

Answers to frequently asked questions about BLENREP



This brochure can support you on your BLENREP treatment journey. It contains answers to the most commonly asked questions about BLENREP. If you have additional questions, speak with your oncologist and healthcare team—they are there for you.

Approved Use

BLENREP is a prescription medicine used in combination with the medicines bortezomib and dexamethasone to treat adults with multiple myeloma who:

- have received at least 2 prior treatments, including a proteasome inhibitor and an immunomodulatory agent, and
- their cancer has come back or did not respond to prior treatment

It is not known if BLENREP is safe and effective in children.

Important Safety Information

What is the most important information I should know about BLENREP?

BLENREP can cause serious side effects, including:

- **Eye problems.** Eye problems are common with BLENREP and can also be severe. BLENREP can cause changes to the surface of your eye, which can lead to symptoms of eye problems, including:
 - decreased vision
 - blurred vision
 - dry eyes
 - sensitivity to light
 - feeling like something is in your eyes
 - eye irritation
 - eye pain

Please see additional Important Safety Information throughout. Click to see the full [Prescribing Information](#), including [Boxed Warning](#) and [Medication Guide](#) for BLENREP.



BLENREP
belantamab
mafodotin-blmf
for injection 70 mg

FREQUENTLY ASKED QUESTIONS

ABOUT MULTIPLE MYELOMA

What are some key facts about multiple myeloma?

Multiple myeloma is the 2nd most common form of blood cancer. This type of cancer is incurable, but it can be treated. It is common for people with multiple myeloma to have periods of remission. Also, some people's multiple myeloma no longer responds to therapy or their cancer progresses.

Important Safety Information for BLENREP (cont'd)

Ulcers on the surface of the eye (corneal ulcers), including with infection, may also happen during treatment with BLENREP.

Tell your healthcare provider right away if you notice any new or worsening eye symptoms or vision changes during treatment with BLENREP.

Your healthcare provider will refer you to an eye care specialist (such as an ophthalmologist or optometrist) to check your eyes before you start treatment, before you receive each dose of BLENREP, and as needed for any new or worsening eye problems. **It is important that you:**

- Keep all of your eye care appointments because some changes can happen without symptoms and may only be seen on an eye exam
- Use preservative-free artificial tears at least 4 times per day starting with your first infusion and continuing until the end of treatment with BLENREP



ABOUT BLENREP

What is BLENREP?

BLENREP is a prescription medicine that's FDA-approved in combination with the medicines bortezomib and dexamethasone to treat adults with multiple myeloma who:

- have received at least 2 prior treatments, including a proteasome inhibitor and an immunomodulatory agent, and
- their cancer has come back or did not respond to prior treatment

It is not known if BLENREP is safe and effective in children.

How does BLENREP work?

BLENREP is a targeted treatment that binds to a protein found on the surface of myeloma cells called BCMA. BLENREP kills myeloma cells by delivering the anti-cancer medicine into the cell, and then triggers the immune system as another way to kill multiple myeloma cells. BCMA is found at higher levels on myeloma cells. BLENREP can affect normal cells. [Watch how BLENREP works here.](#)

How was BLENREP studied?


BLENREP was given in combination with bortezomib and dexamethasone in a global, randomized clinical study called DREAMM-7.

DREAMM-7 included people with multiple myeloma who had previously been given 2 or more treatments, including a proteasome inhibitor and an immunomodulatory agent, and experienced disease progression while taking, or after, their most recent treatment.

Important Safety Information for BLENREP (cont'd)

- Avoid wearing contact lenses during treatment with BLENREP unless directed by your eye care specialist
- Use caution when driving or operating machinery because BLENREP may cause changes to your vision

Please see additional Important Safety Information throughout. Click to see the full Prescribing Information, including Boxed Warning and Medication Guide for BLENREP.

 **BLENREP**
belantamab
mafodotin-blmf
for injection 70 mg

FREQUENTLY ASKED QUESTIONS



ABOUT BLENREP (CONT'D)

How will I receive BLENREP?

BLENREP will be given to you through an intravenous (IV) infusion at a Risk Evaluation and Mitigation Strategy (REMS)-certified outpatient site, such as your doctor's office or an outpatient clinic. Timing will vary by patient, but can be as short as 30 minutes. No hospitalization is needed to start treatment with BLENREP.

- BLENREP is used in combination with bortezomib and dexamethasone. Dosing schedule usually starts with infusions every 3 weeks for the first 8 cycles, then continues with BLENREP alone
- Your healthcare provider will decide how many treatments you will need
- Your healthcare provider will check your blood cell counts before you start treatment with BLENREP, and during treatment as needed
- If you have side effects, your oncologist may adjust your dose. This may include an increase in time between doses or a decrease in dose
- If you experience serious side effects, your oncologist may temporarily or completely stop treatment with BLENREP

TAKING BLENREP

Why is BLENREP only available through REMS?

Because of the risk of eye problems, BLENREP is available only through a restricted program called the BLENREP Risk Evaluation and Mitigation Strategy (REMS).

- REMS helps your care team monitor your eye health
- You will receive the BLENREP REMS Patient Guide from your care team
- You will receive counseling on the risk of eye problems and the need for eye exams
- To receive BLENREP, you must enroll in the REMS program and receive eye exams during treatment

[Click here to learn more about the BLENREP REMS program and talk to your doctor.](#)

Important Safety Information for BLENREP (cont'd)

Because of the risk of eye problems, BLENREP is available only through a restricted program called the BLENREP Risk Evaluation and Mitigation Strategy (REMS). Your healthcare provider will give you the BLENREP REMS Patient Guide, which describes the risk of eye problems, and will explain the REMS program to you. To receive BLENREP, you must enroll in the REMS program and receive eye exams during treatment. See **"What are the possible side effects of BLENREP?"** for more information about side effects.

TAKING BLENREP (CONT'D)

Why do I need to get my eyes checked?

Eye problems are common with BLENREP and may also be severe. BLENREP can cause changes to the surface of your eye, which may result in changes in vision and symptoms of eye problems, such as:

- decreased vision
- blurred vision
- dry eyes
- sensitivity to light
- feeling like something is in your eyes
- eye irritation
- eye pain

Also, ulcers on the surface of the eye, known as corneal ulcers, including with infection, may happen during treatment with BLENREP.

Even if your vision seems fine, it is important that you get your eyes checked during treatment with BLENREP because some changes can happen without symptoms, and may only be seen on an eye exam. It is important to tell your healthcare provider if you notice any new or worsening eye symptoms during treatment with BLENREP.

What do I need to do before starting BLENREP?

Eye exams are required before starting BLENREP, then repeated before each dose. Additional eye exams will also be needed for new or worsening eye problems or as your doctor recommends. This is done to make sure your healthcare team can monitor any changes that occur and determine if a change in dose is needed.


Eye exams include:

- Best-Corrected Visual Acuity test, where you read an eye chart from a distance
- Slit lamp test, where an eye doctor takes a closer look at the front and inside of your eye

Important Safety Information for BLENREP (cont'd)

What are the possible side effects of BLENREP?
BLENREP can cause serious side effects, including:
• See **"Eye Problems"** in **"What is the most important information I should know about BLENREP?"**

Please see additional Important Safety Information throughout. Click to see the full [Prescribing Information](#), including [Boxed Warning](#) and [Medication Guide](#) for BLENREP.

 **BLENREP**
belantamab
mafodotin-blmf
for injection 70 mg

FREQUENTLY ASKED QUESTIONS

TAKING BLENREP (CONT'D)

Are there any tips to help with eye problems?

Keep all your eye care appointments because some changes can happen without symptoms and may only be seen on an eye exam.

It is important that you use preservative-free artificial tears at least 4 times per day during treatment with BLENREP to help manage symptoms.

You should avoid wearing contact lenses during treatment with BLENREP, unless you are directed to by an eye doctor (an ophthalmologist or optometrist).

Are there any restrictions on certain daily activities involving vision after starting BLENREP?

BLENREP may cause changes to your vision. Use caution when driving or operating machinery. Talk to your oncologist if you are not sure.

In addition to eye problems, are there other serious side effects that can occur with BLENREP?

Some serious side effects that can occur while taking BLENREP include the following:

- BLENREP can also cause a decrease in platelets (thrombocytopenia), which is common with BLENREP and can also be severe. Platelets are a type of blood cell that help your blood to clot. Your healthcare provider will check your blood cell counts before you start treatment with BLENREP and during treatment as needed. Tell your healthcare provider if you get any bleeding or bruising during treatment with BLENREP



TAKING BLENREP (CONT'D)

What are the most common side effects with BLENREP?

In addition to eye problems, the most common side effects of BLENREP when given with bortezomib and dexamethasone include:

- upper respiratory tract infection
- liver problems
- diarrhea
- tiredness
- pneumonia
- cataract
- COVID-19

The most common severe abnormal blood test results during treatment with BLENREP include decreases in platelets, white blood cells, and hemoglobin, and increases in certain liver enzymes.

Tell your healthcare provider right away if you get new or worsening unexplained signs or symptoms of lung problems, including shortness of breath, chest pain, or cough.

BLENREP may affect fertility in males and females, which may affect your ability to have children. Talk to your healthcare provider if this is a concern for you.


These are not all the possible side effects of BLENREP.

Will I have to stop my treatment if I experience eye problems?

- If you have side effects, your oncologist may adjust your dose
- Adjustments may include an increase in time between doses or a decrease in dose
- If you experience serious side effects, your oncologist may temporarily or completely stop treatment with BLENREP
- If you miss any appointments, call your healthcare provider as soon as possible to reschedule your appointment

Talk to your doctor for medical advice about side effects.

Please see Important Safety Information throughout. Click to see the full [Prescribing Information](#), including [Boxed Warning](#) and [Medication Guide](#) for BLENREP.

 **BLENREP**
belantamab
mafodotin-blmf
for injection 70 mg

FREQUENTLY ASKED QUESTIONS

TAKING BLENREP (CONT'D)

Could BLENREP affect my vision?

Yes, eye problems are common with BLENREP. BLENREP can cause changes to the surface of your eye, which may result in changes to your vision. Symptoms such as decreased vision, blurred vision, dry eyes, sensitivity to light, feeling like something is in your eyes, eye irritation, or eye pain are common.

Severe reduction in vision or ulcers on the surface of the eye (corneal ulcers), including with infection, may happen during treatment with BLENREP. It is important to tell your healthcare provider right away if you notice any new or worsening eye symptoms, or changes in vision during treatment with BLENREP. Your healthcare provider may decrease your dose, temporarily stop, or completely stop treatment with BLENREP if you get serious side effects. During the clinical trial, 83% of patients required a dosage modification due to eye problems.

What should I do if I can't make it to an eye exam?

Contact your eye care professional or nurse, who will help you reschedule your eye exam as soon as possible, so you can stay on treatment. It is important to work with your eye care professional to schedule your appointment to avoid delays in getting your dose of BLENREP.

Who do I need to see to get my eyes checked?

Optometrists are eye care professionals who are qualified to perform eye exams but are not medical doctors. Ophthalmologists are medical doctors who specialize in eyes and eye care. You can visit either of these eye care professionals for the eye exams that are needed for BLENREP treatment.

Important Safety Information for BLENREP (cont'd)

- **Decrease in platelets (thrombocytopenia)** is common with BLENREP and can also be severe. Platelets are a type of blood cell that help your blood to clot. Your healthcare provider will check your blood cell counts before you start treatment with BLENREP and during treatment as needed. Tell your healthcare provider if you get any bleeding or bruising during treatment with BLENREP

The most common side effects of BLENREP when given with bortezomib and dexamethasone include:

- See "Eye problems" in "What is the most important information I should know about BLENREP?"
- upper respiratory tract infection
- liver problems
- diarrhea
- tiredness
- pneumonia
- cataract
- COVID-19

SUPPORT AND RESOURCES



What is Together with BLENREP?

Together with BLENREP is designed to support patients like you during treatment with BLENREP. It is more than a program—it's our promise to understand your needs, listen to what's important to you, and create meaningful connections along the way.

For patients prescribed BLENREP, Together with BLENREP offers personalized help to get you started and stay informed. For more information on program features, visit togetherwithBLENREP.com.

How can I find out if eye exams are covered under insurance?

You can contact your insurer directly to determine whether your eye exams will be covered. If you sign up for Together with BLENREP, your dedicated Nurse Navigator will have your insurance coverage checked and help you understand your benefits. The information provided by Together with BLENREP is not a guarantee of coverage or payment.

What is a Together with BLENREP Nurse Navigator?

Your dedicated Nurse Navigator is someone who can help you, your oncologist's office, and your eye care professional (ECP)'s office work together during your treatment journey with BLENREP.

Keep in mind that the Nurse Navigator does not provide medical advice and is not a part of your doctor's care team. Please ask your doctor any questions that you might have about your disease and treatment with BLENREP.


Important Safety Information for BLENREP (cont'd)

The most common severe abnormal blood test results during treatment with BLENREP include decreases in platelets, white blood cells, and hemoglobin, and increases in certain liver enzymes.

Tell your healthcare provider right away if you get new or worsening unexplained signs or symptoms of lung problems, including shortness of breath, chest pain, or cough.

Your healthcare provider may decrease your dose, temporarily stop, or completely stop treatment with BLENREP if you get serious side effects.

Please see additional Important Safety Information throughout. Click to see the full Prescribing Information, including Boxed Warning and Medication Guide for BLENREP.

 **BLENREP**
belantamab
mafodotin-blmf
for injection 70 mg

Important Safety Information for BLENREP

What is the most important information I should know about BLENREP?

BLENREP can cause serious side effects, including:

- **Eye problems.** Eye problems are common with BLENREP and can also be severe. BLENREP can cause changes to the surface of your eye, which can lead to symptoms of eye problems, including:
 - decreased vision
 - blurred vision
 - dry eyes
 - sensitivity to light
 - feeling like something is in your eyes
 - eye irritation
 - eye pain

Ulcers on the surface of the eye (corneal ulcers), including with infection, may also happen during treatment with BLENREP.

Tell your healthcare provider right away if you notice any new or worsening eye symptoms or vision changes during treatment with BLENREP.

Your healthcare provider will refer you to an eye care specialist (such as an ophthalmologist or optometrist) to check your eyes before you start treatment, before you receive each dose of BLENREP, and as needed for any new or worsening eye problems. **It is important that you:**

- Keep all of your eye care appointments because some changes can happen without symptoms and may only be seen on an eye exam
- Use preservative-free artificial tears at least 4 times per day starting with your first infusion and continuing until the end of treatment with BLENREP
- Avoid wearing contact lenses during treatment with BLENREP unless directed by your eye care specialist
- Use caution when driving or operating machinery because BLENREP may cause changes to your vision

Because of the risk of eye problems, BLENREP is available only through a restricted program called the BLENREP Risk Evaluation and Mitigation Strategy (REMS). Your healthcare provider will give you the BLENREP REMS Patient Guide, which describes the risk of eye problems, and will explain the REMS program to you. To receive BLENREP, you must enroll in the REMS program and receive eye exams during treatment. See **“What are the possible side effects of BLENREP?”** for more information about side effects.

What are the possible side effects of BLENREP?

BLENREP can cause serious side effects, including:

- See **“Eye Problems”** in **“What is the most important information I should know about BLENREP?”**
- **Decrease in platelets (thrombocytopenia)** is common with BLENREP and can also be severe. Platelets are a type of blood cell that help your blood to clot. Your healthcare provider will check your blood cell counts before you start treatment with BLENREP and during treatment as needed. Tell your healthcare provider if you get any bleeding or bruising during treatment with BLENREP

The most common side effects of BLENREP when given with bortezomib and dexamethasone include:

- See **“Eye problems”** in **“What is the most important information I should know about BLENREP?”**
- upper respiratory tract infection
- liver problems
- diarrhea
- tiredness
- pneumonia
- cataract
- COVID-19

The most common severe abnormal blood test results during treatment with BLENREP include decreases in platelets, white blood cells, and hemoglobin, and increases in certain liver enzymes.

Tell your healthcare provider right away if you get new or worsening unexplained signs or symptoms of lung problems, including shortness of breath, chest pain, or cough.

Your healthcare provider may decrease your dose, temporarily stop, or completely stop treatment with BLENREP if you get serious side effects.

BLENREP may affect fertility in males and females, which may affect your ability to have children. Talk to your healthcare provider if this is a concern for you.

These are not all of the possible side effects of BLENREP.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Before receiving BLENREP, tell your healthcare provider about all of your medical conditions, including if you:

- have or had vision or eye problems
- have or had bleeding problems
- are pregnant or plan to become pregnant. BLENREP can harm your unborn baby

Females who are able to become pregnant:


- Your healthcare provider will do a pregnancy test before you start treatment with BLENREP
- Use effective birth control during treatment with BLENREP and for 4 months after the last dose. Talk to your healthcare provider about birth control methods you can use during this time
- Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment with BLENREP

Males with female partners who are able to become pregnant:

- Use effective birth control during treatment with BLENREP and for 6 months after the last dose
- are breastfeeding or plan to breastfeed. It is not known if BLENREP passes into your breast milk. Do not breastfeed during treatment with BLENREP and for 3 months after the last dose

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088. You may also report negative side effects to GSK at gsk.public.reportum.com or 1-888-825-5249.



Please click to see the full [Prescribing Information](#), including [Boxed Warning](#) and [Medication Guide](#) for BLENREP.

If you haven't enrolled in Together with BLENREP, sign up today!



Click the link below to enroll and access additional resources, Nurse Navigators,* and much more.



We are here to help!

You can reach your Nurse Navigator at **1-844-4GSK-ONC** (1-844-447-5662)

Monday–Friday 8:00 AM–8:00 PM ET
Or visit togetherwithBLENREP.com

*The Nurse Navigator does not provide medical advice and is not a part of your doctor's care team. Please ask your doctor any questions that you might have about your disease and treatment with BLENREP.

Please see Important Safety Information throughout. Click to see the full Prescribing Information, including Boxed Warning and Medication Guide for BLENREP.

Trademarks are owned by or licensed to the GSK group of companies.



©2025 GSK or licensor.
PMUS-BLMBKLT250041 December 2025
Produced in USA.



BLENREP
belantamab
mafodotin-blmf
for injection 70 mg