

BLENREP
belantamab
mafodotin-blmf
for injection 70 mg

CARE PARTNER GUIDE

I want to be a
come

**back
kid**

Hilary

**BLENREP may help power
a comeback to remission if her
multiple myeloma returns**

Not an actual patient
or caregiver.

HELP SUPPORT YOUR LOVED ONE IF MULTIPLE MYELOMA RETURNS

Use this brochure to find out more about BLENREP, how you can be there for your loved one throughout their treatment, and find available support and resources.*

*This information is not a substitute for a healthcare provider's advice. Your loved one's care team is the best source for answers to questions about their treatment plan.

Approved Use

BLENREP is a prescription medicine used in combination with the medicines bortezomib and dexamethasone to treat adults with multiple myeloma who:

- have received at least 2 prior treatments, including a proteasome inhibitor and an immunomodulatory agent, and
- their cancer has come back or did not respond to prior treatment

It is not known if BLENREP is safe and effective in children.

Important Safety Information

What is the most important information I should know about BLENREP?

BLENREP can cause serious side effects, including:

- **Eye problems.** Eye problems are common with BLENREP and can also be severe. BLENREP can cause changes to the surface of your eye, which can lead to symptoms of eye problems, including:
 - decreased vision
 - blurred vision
 - dry eyes
 - sensitivity to light
 - feeling like something is in your eyes
 - eye irritation
 - eye pain

Please see additional Important Safety Information throughout. Click to see the full [Prescribing Information](#), including [Boxed Warning](#) and [Medication Guide](#) for BLENREP.

What is BLENREP?



BLENREP is a prescription medicine that's FDA-approved for use in combination with the medicines bortezomib and dexamethasone to treat adults with multiple myeloma:

- who have received at least 2 prior treatments, including a proteasome inhibitor and an immunomodulatory agent, and
- whose cancer has come back or did not respond to prior treatment

It is not known if BLENREP is safe and effective in children.

Christopher

BLENREP may help power a comeback to remission if his multiple myeloma returns

Not an actual patient or caregiver.

Important Safety Information for BLENREP (cont'd)

What is the most important information I should know about BLENREP? (cont'd)

BLENREP can cause serious side effects, including:

Ulcers on the surface of the eye (corneal ulcers), including with infection, may also happen during treatment with BLENREP.

Tell your healthcare provider right away if you notice any new or worsening eye symptoms or vision changes during treatment with BLENREP.

You, your loved one, and treatment with BLENREP

Being a care partner means helping your loved one as they go through multiple myeloma treatment and supporting their overall wellness

It takes time, energy, and effort to be a care partner—go to the appointments, make the calls, and help manage their day-to-day needs.

How this brochure can help support you



Use this brochure to find key information about BLENREP. You'll find answers to some questions that you may have like:

- What are the steps in the treatment journey with BLENREP?
- What are the possible side effects of BLENREP?
- What support and resources are available to me and my loved one while they are given BLENREP?

Care teams, care partners, and patients all work toward the common goal of getting multiple myeloma in remission, and BLENREP may be able to help.

Remission: Refers to progression-free survival, the average length of time after the start of treatment in which a person is alive and their cancer does not get worse.

Important Safety Information for BLENREP (cont'd)

What is the most important information I should know about BLENREP? (cont'd)

Your healthcare provider will refer you to an eye care specialist (such as an ophthalmologist or optometrist) to check your eyes before you start treatment, before you receive each dose of BLENREP, and as needed for any new or worsening eye problems. **It is important that you:**

- Keep all of your eye care appointments because some changes can happen without symptoms and may only be seen on an eye exam
- Use preservative-free artificial tears at least 4 times per day starting with your first infusion and continuing until the end of treatment with BLENREP
- Avoid wearing contact lenses during treatment with BLENREP unless directed by your eye care specialist
- Use caution when driving or operating machinery because BLENREP may cause changes to your vision

Please see additional Important Safety Information throughout. Click to see the full Prescribing Information, including Boxed Warning and Medication Guide for BLENREP.

BLENREP
belantamab
mafodotin-blmf
for injection 70 mg


BLENREP is a targeted treatment that works in multiple ways

How BLENREP works

BLENREP binds to a protein found on the surface of myeloma cells called BCMA

BLENREP KILLS MYELOMA CELLS

by delivering the anti-cancer medicine into the cell



BLENREP TRIGGERS THE IMMUNE SYSTEM

as another way to kill multiple myeloma cells

BCMA is found at higher levels on myeloma cells. BLENREP can affect normal cells.

Maria

BLENREP may help power a comeback to remission if her multiple myeloma returns

Not an actual patient or caregiver.

Important Safety Information for BLENREP (cont'd)

What is the most important information I should know about BLENREP? (cont'd)

Because of the risk of eye problems, BLENREP is available only through a restricted program called the BLENREP Risk Evaluation and Mitigation Strategy (REMS). Your healthcare provider will give you the BLENREP REMS Patient Guide, which describes the risk of eye problems, and will explain the REMS program to you. To receive BLENREP, you must enroll in the REMS program and receive eye exams during treatment. See **“What are the possible side effects of BLENREP?”** for more information about side effects.

What to expect when you start BLENREP

How is BLENREP given



BLENREP is given as an intravenous (IV) infusion in a **Risk Evaluation and Mitigation Strategy (REMS)**-certified outpatient site, such as a doctor's office or an outpatient clinic. Your loved one must be enrolled in the REMS program and receive eye exams before each dose for as long as they're on treatment.

The dosing schedule usually starts with infusions of BLENREP every 3 weeks, used in combination with bortezomib and dexamethasone for the first 8 cycles. Then treatment continues with BLENREP alone.

Timing will vary by patient, but infusions of BLENREP can be as short as 30 minutes.

No hospitalization is needed to start treatment with BLENREP.

REMS: Program that helps your loved one's care team monitor their eye health and determine if the benefits of BLENREP outweigh the risks.

Important Safety Information for BLENREP (cont'd)

What are the possible side effects of BLENREP?

BLENREP can cause serious side effects, including:

- See **“Eye Problems”** in **“What is the most important information I should know about BLENREP?”**
- **Decrease in platelets (thrombocytopenia)** is common with BLENREP and can also be severe. Platelets are a type of blood cell that help your blood to clot. Your healthcare provider will check your blood cell counts before you start treatment with BLENREP and during treatment as needed. Tell your healthcare provider if you get any bleeding or bruising during treatment with BLENREP

Please see additional Important Safety Information throughout. Click to see the full Prescribing Information, including Boxed Warning and Medication Guide for BLENREP.

BLENREP
belantamab
mafodotin-blmf
for injection 70 mg

Guide to the BLENREP treatment journey

1 Review any informational materials provided by your healthcare team if your loved one is prescribed BLENREP.

2 Your loved one enrolls in the mandatory REMS program and can choose to enroll in the Together with BLENREP Program.

Once enrolled in REMS, your loved one will receive the BLENREP REMS Patient Guide from their care team, which will explain the risk of eye problems. If they choose to enroll in Together with BLENREP, they can access additional resources, Nurse Navigators, and much more. For more information on how to enroll, see pages 10 and 11 of this guide.

3 Help your loved one find an eye doctor.

Having an eye doctor is needed since eye exams are required before treatment starts and then before each dose of BLENREP. Additional eye exams will also be needed for new or worsening symptoms or as their doctor recommends. This is because eye problems such as changes in vision are common with BLENREP treatment and may only be seen on an eye exam.



The oncologist may also refer your loved one to an eye doctor, or you both can visit [BLENREP.com](https://www.blenrep.com) and use the "Find an Eye Doctor" tool to locate one near by.

4 Schedule an eye exam appointment.

If your loved one misses any appointments, it is important to work with their care team to reschedule as soon as possible. This is to avoid delays in their next dose of BLENREP.

5 Help your loved one schedule their BLENREP infusion once their oncologist receives the eye exam results.

Keep track of the dates of BLENREP infusions and adjust appointments if needed.

6 Your loved one will be given their BLENREP infusion.

If your loved one experiences side effects throughout treatment, their oncologist may increase time between doses, adjust their dose, or temporarily or completely stop treatment with BLENREP.

On treatment days, you can help by packing items that may give your loved one comfort, such as a blanket or heating pad. You can also be there to ask the care team questions when needed.



Repeat steps 4–6 for each dose.

7 Help support your loved one in discussing their experience on treatment with their oncologist, including any side effects they may experience.

Keep track of important information related to treatment in a journal or notebook. This can inform your loved one's oncologist of experiences between appointments. Your loved one will be told to use preservative-free artificial tears at least 4 times per day starting with their first infusion and continuing until the end of treatment with BLENREP.

Important Safety Information for BLENREP (cont'd)

What are the possible side effects of BLENREP? (cont'd)

The most common side effects of BLENREP when given with bortezomib and dexamethasone include:


- See "Eye problems" in "What is the most important information I should know about BLENREP?"
- upper respiratory tract infection
- diarrhea
- pneumonia
- COVID-19
- liver problems
- tiredness
- cataract

Important Safety Information for BLENREP (cont'd)

What are the possible side effects of BLENREP? (cont'd)

The most common severe abnormal blood test results during treatment with BLENREP include decreases in platelets, white blood cells, and hemoglobin, and increases in certain liver enzymes.

Please see additional Important Safety Information throughout. Click to see the full [Prescribing Information](#), including [Boxed Warning](#) and [Medication Guide](#) for BLENREP.



The most important information that you and your loved one should know about BLENREP

BLENREP can cause serious side effects, including:

- Eye problems, which are common with BLENREP and can also be severe. BLENREP can cause changes to the surface of your eye, which can lead to symptoms of eye problems, including:
 - decreased vision
 - blurred vision
 - dry eyes
 - sensitivity to light
 - feeling like something is in your eyes
 - eye irritation
 - eye pain

Ulcers on the surface of the eye (corneal ulcers), including with infection, may also happen during treatment with BLENREP.

Tell your healthcare provider right away if you notice any new or worsening eye symptoms or vision changes during treatment with BLENREP.

Your healthcare provider will refer you to an eye doctor (such as an ophthalmologist or optometrist) to check your eyes before you start treatment, before you receive each dose of BLENREP, and as needed for any new or worsening eye problems. **It is important that you:**

- Keep all of your eye care appointments because some changes can happen without symptoms and may only be seen on an eye exam
- Use preservative-free artificial tears at least 4 times per day starting with your first infusion and continuing until the end of treatment with BLENREP
- Avoid wearing contact lenses during treatment with BLENREP unless directed by their eye doctor
- Use caution when driving or operating machinery because BLENREP may cause changes to their vision

Because of the risk of eye problems, BLENREP is available only through a restricted program called the BLENREP Risk Evaluation and Mitigation Strategy (REMS). Your healthcare provider will give you the BLENREP REMS Patient Guide, which describes the risk of eye problems, and will explain the REMS program to you. To receive BLENREP, you must enroll in the REMS program and receive eye exams during treatment.



In addition to eye problems, it is important to be aware of other side effects that may occur

Additional serious side effects can include

- **Decrease in platelets (thrombocytopenia)** is common with BLENREP and can also be severe. Platelets are a type of blood cell that help your blood to clot. Your healthcare provider will check your blood cell counts before you start treatment with BLENREP and during treatment as needed. Tell your healthcare provider if you get any bleeding or bruising during treatment with BLENREP

In addition to eye problems, the most common side effects of BLENREP when given with bortezomib and dexamethasone include:

- upper respiratory tract infection
- liver problems
- diarrhea
- tiredness
- pneumonia
- cataract
- COVID-19

The most common severe abnormal blood test results during treatment with BLENREP include decreases in platelets, white blood cells, and hemoglobin, and increases in certain liver enzymes.

Tell your healthcare provider right away if you get new or worsening unexplained signs or symptoms of lung problems, including shortness of breath, chest pain, or cough.

Your healthcare provider may decrease your dose, temporarily stop, or completely stop treatment with BLENREP if you get serious side effects.

BLENREP may affect fertility in males and females, which may affect your ability to have children. Talk to your healthcare provider if this is a concern for you.

These are not all of the possible side effects of BLENREP.

Please see additional Important Safety Information throughout. Click to see the full [Prescribing Information](#), including [Boxed Warning](#) and [Medication Guide](#) for BLENREP.

The Together with BLENREP program is support made for you and your loved one

Together with BLENREP is designed to support patients and their care partners during treatment with BLENREP.

It's more than a program—it's our promise to understand your needs, listen to what's important to you, and create meaningful connections along the way.

For patients prescribed BLENREP, Together with BLENREP offers personalized help to get them started and stay informed.

Key program features include:

- Coverage support: Insurance and coverage information to your loved one and their doctor, including supporting their doctor if they face coverage challenges
- Cost support: Help make paying for BLENREP more manageable with options tailored to your loved one's needs
- Patient Support: A dedicated Nurse Navigator* through key steps in your loved one's treatment journey by helping them:
 - Understand their insurance coverage
 - Find an eye care professional (ECP) for required eye exams
 - Receive education about BLENREP and multiple myeloma

You can help them enroll in 2 ways:

- Visit www.togetherwithGSK.com/blenrep/patient/ to start enrollment online; or
- Enroll by downloading this form and work with your loved one's doctor to fill it out at their visit for their doctor to submit

*The Nurse Navigator does not provide medical advice and is not a part of your doctor's care team.

If your loved one hasn't enrolled in Together with BLENREP, help them sign up today! [Click here.](#)



Resources you both can access online

Online resources for other organizations can be found at:

- [Blood Cancer United](#)
- [CancerCare](#)
- [Cancer Support Community](#)
- [HealthTree](#)
- [International Myeloma Foundation](#)
- [Multiple Myeloma Research Foundation](#)

Please note: These resources are external to GSK. GSK does not have control over the content or information provided through these resources, and accordingly does not warrant their accuracy or completeness. These organizations do not endorse any products or services.

Educational resources about BLENREP are available at BLENREP.com



We are here to help!

Once enrolled in Together with BLENREP, your loved one can reach their Nurse Navigator* at **1-844-4GSK-ONC** (1-844-447-5662) Monday–Friday 8:00 am–8:00 pm ET. Or visit togetherwithBLENREP.com

*The Nurse Navigator does not provide medical advice and is not a part of your doctor's care team. Please ask your doctor any questions that you might have about your disease and treatment with BLENREP.

Henry

BLENREP may help power a comeback to remission if his multiple myeloma returns

Not an actual patient or caregiver.



Please see Important Safety Information throughout. Click to see the full [Prescribing Information](#), including [Boxed Warning](#) and [Medication Guide](#) for BLENREP.

BLENREP
belantamab
mafodotin-blmf
for injection 70 mg

How to be a voice in the room



- With your loved one's permission, talk with the care team to help align the care plan with your loved one's goals and share updates as goals change over time
- You can also be the one who listens and remembers what is being explained to your loved one
- When with the care team, speak up if you don't understand something to help get the information you both need
- The care team may not know what your loved one is feeling until they are told

As a care partner, you can offer support to help your loved one and help improve communication with their care team

Hilary

BLNREP may help power a comeback to remission if her multiple myeloma returns

Not an actual patient or caregiver.



Important Safety Information for BLNREP

What is the most important information I should know about BLNREP?

BLNREP can cause serious side effects, including:

- **Eye problems.** Eye problems are common with BLNREP and can also be severe. BLNREP can cause changes to the surface of your eye, which can lead to symptoms of eye problems, including:
 - decreased vision
 - blurred vision
 - dry eyes
 - sensitivity to light
 - feeling like something is in your eyes
 - eye irritation
 - eye pain

Ulcers on the surface of the eye (corneal ulcers), including with infection, may also happen during treatment with BLNREP.

Tell your healthcare provider right away if you notice any new or worsening eye symptoms or vision changes during treatment with BLNREP.

Your healthcare provider will refer you to an eye care specialist (such as an ophthalmologist or optometrist) to check your eyes before you start treatment, before you receive each dose of BLNREP, and as needed for any new or worsening eye problems. **It is important that you:**

- Keep all of your eye care appointments because some changes can happen without symptoms and may only be seen on an eye exam
- Use preservative-free artificial tears at least 4 times per day starting with your first infusion and continuing until the end of treatment with BLNREP
- Avoid wearing contact lenses during treatment with BLNREP unless directed by your eye care specialist
- Use caution when driving or operating machinery because BLNREP may cause changes to your vision

Because of the risk of eye problems, BLNREP is available only through a restricted program called the BLNREP Risk Evaluation and Mitigation Strategy (REMS). Your healthcare provider will give you the BLNREP REMS Patient Guide, which describes the risk of eye problems, and will explain the REMS program to you. To receive BLNREP, you must enroll in the REMS program and receive eye exams during treatment. See **"What are the possible side effects of BLNREP?"** for more information about side effects.

Please click to see the full [Prescribing Information](#), including [Boxed Warning](#) and [Medication Guide](#) for BLNREP.

BLNREP
belantamab
mafodotin-blmf
for injection 70 mg

Important Safety Information for BLENREP (cont'd)

What are the possible side effects of BLENREP?

BLENREP can cause serious side effects, including:

- See “Eye Problems” in “What is the most important information I should know about BLENREP?”
- **Decrease in platelets (thrombocytopenia)** is common with BLENREP and can also be severe. Platelets are a type of blood cell that help your blood to clot. Your healthcare provider will check your blood cell counts before you start treatment with BLENREP and during treatment as needed. Tell your healthcare provider if you get any bleeding or bruising during treatment with BLENREP

The most common side effects of BLENREP when given with bortezomib and dexamethasone include:

- See “Eye problems” in “What is the most important information I should know about BLENREP?”
- upper respiratory tract infection
- liver problems
- diarrhea
- tiredness
- pneumonia
- cataract
- COVID-19

The most common severe abnormal blood test results during treatment with BLENREP include decreases in platelets, white blood cells, and hemoglobin, and increases in certain liver enzymes.

Tell your healthcare provider right away if you get new or worsening unexplained signs or symptoms of lung problems, including shortness of breath, chest pain, or cough.

Your healthcare provider may decrease your dose, temporarily stop, or completely stop treatment with BLENREP if you get serious side effects.

BLENREP may affect fertility in males and females, which may affect your ability to have children. Talk to your healthcare provider if this is a concern for you.

These are not all of the possible side effects of BLENREP.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Before receiving BLENREP, tell your healthcare provider about all of your medical conditions, including if you:

- have or had vision or eye problems
- have or had bleeding problems
- are pregnant or plan to become pregnant. BLENREP can harm your unborn baby

Females who are able to become pregnant:

- Your healthcare provider will do a pregnancy test before you start treatment with BLENREP
- Use effective birth control during treatment with BLENREP and for 4 months after the last dose. Talk to your healthcare provider about birth control methods you can use during this time
- Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment with BLENREP

Males with female partners who are able to become pregnant:

- Use effective birth control during treatment with BLENREP and for 6 months after the last dose
- are breastfeeding or plan to breastfeed. It is not known if BLENREP passes into your breast milk. Do not breastfeed during treatment with BLENREP and for 3 months after the last dose

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088. You may also report negative side effects to GSK at gsk.public.reportum.com or 1-888-825-5249.




Please click to see the full [Prescribing Information](#), including [Boxed Warning](#) and [Medication Guide](#) for BLENREP.

BLENREP
belantamab
mafodotin-blmf
for injection 70 mg

I want to be a

come back kid

CARE PARTNER GUIDE



Henry
BLENREP may help power
a comeback to remission if his
multiple myeloma returns

Not an actual patient
or caregiver.

**HELP SUPPORT YOUR LOVED
ONE ON THEIR TREATMENT
JOURNEY WITH BLENREP**

Approved Use

BLENREP is a prescription medicine used in combination with the medicines bortezomib and dexamethasone to treat adults with multiple myeloma who:

- have received at least 2 prior treatments, including a proteasome inhibitor and an immunomodulatory agent, and
- their cancer has come back or did not respond to prior treatment

It is not known if BLENREP is safe and effective in children.

Please see additional Important Safety Information throughout. Click to see the full [Prescribing Information](#), including [Boxed Warning](#) and [Medication Guide](#) for BLENREP.

Important Safety Information

What is the most important information I should know about BLENREP?

BLENREP can cause serious side effects, including:

- **Eye problems.** Eye problems are common with BLENREP and can also be severe. BLENREP can cause changes to the surface of your eye, which can lead to symptoms of eye problems, including:
 - decreased vision
 - blurred vision
 - dry eyes
 - sensitivity to light
 - feeling like something is in your eyes
 - eye irritation
 - eye pain

Trademarks are owned by or licensed to the GSK group of companies.

GSK

©2026 GSK or licensor.
PMUS-BLMLBND250013 January 2026
Produced in USA.

File Name: Care Partner Guide



BLENREP
belantamab
mafodotin-blmf
for injection 70 mg