



Not an actual patient
or care partner.

**BLENREP may help
power a comeback to
remission* if her multiple
myeloma returns**

YOUR ROLE IN THE JOURNEY MATTERS

A care partner's guide to supporting treatment for loved ones prescribed BLENREP

This information is not a substitute for a healthcare provider's advice. Your loved one's care team is the best source for answers to questions about their treatment plan.

*Remission here means median progression-free survival, the length of time after the start of treatment in which a person is alive and their cancer does not get worse.

What is BLENREP?

BLENREP is a prescription medicine used in combination with the medicines bortezomib and dexamethasone to treat adults with multiple myeloma who:

- have received at least 2 prior treatments, including a proteasome inhibitor and an immunomodulatory agent, and
- their cancer has come back or did not respond to prior treatment.

It is not known if BLENREP is safe and effective in children.

Important Safety Information

What is the most important information I should know about BLENREP?

BLENREP can cause serious side effects, including:

- **Eye problems.** Eye problems are common with BLENREP and can also be severe. BLENREP can cause changes to the surface of your eye, which can lead to symptoms of eye problems, including:
 - decreased vision
 - blurred vision
 - dry eyes
 - sensitivity to light
 - feeling like something is in your eyes
 - eye irritation
 - eye pain

Please see additional Important Safety Information throughout. Click to see the full [Prescribing Information](#), including [Boxed Warning](#) and [Medication Guide](#) for BLENREP.

BLENREP
belantamab
mafodotin-blmf
for injection 70 mg

Welcome to the BLENREP journey

Caring for someone with multiple myeloma is one of the most meaningful things a person can do. It's also one of the most demanding. The role of a care partner asks much of your time, your energy, and your emotions.

This guide was created with you in mind—to give you the information, resources, and reassurance you need as you support and care for your loved one on the BLENREP combination.

Guide to the BLENREP treatment journey

Overview of the BLENREP clinical study

Eye and vision changes with BLENREP

Getting started with BLENREP

Helpful reminders for care partners

Support programs and resources

4

6

8

10

11

12

To get a better understanding of how BLENREP works, [click here](#)

Important Safety Information for BLENREP (cont'd)

Ulcers on the surface of the eye (corneal ulcers), including with infection, may also happen during treatment with BLENREP.

Tell your healthcare provider right away if you notice any new or worsening eye symptoms or vision changes during treatment with BLENREP.

Your healthcare provider will refer you to an eye care specialist (such as an ophthalmologist or optometrist) to check your eyes before you start treatment, before you receive each dose of BLENREP, and as needed for any new or worsening eye problems.

This guide was made specifically for care partners like you

“I can’t protect my wife from something that comes from the inside, but I can be there to hold her hand.”

–Real care partner like you

“Sometimes ... it is nice when someone helps you to talk about things that are difficult to talk about.”

–Real care partner like you



We want to help you focus on what’s important



Not an actual patient or care partner.

Important Safety Information for BLENREP (cont'd)

It is important that you:

- Keep all of your eye care appointments because some changes can happen without symptoms and may only be seen on an eye exam
- Use preservative-free artificial tears at least 4 times per day starting with your first infusion and continuing until the end of treatment with BLENREP
- Avoid wearing contact lenses during treatment with BLENREP unless directed by your eye care specialist
- Use caution when driving or operating machinery because BLENREP may cause changes to your vision

Please see additional Important Safety Information throughout. Click to see the full Prescribing Information, including Boxed Warning and Medication Guide for BLENREP.



Guide to the BLENREP treatment journey

Your care team is here to support you and your loved one every step of the way.

1. ENROLL

Because of the risk of eye problems, your loved one must enroll in the mandatory safety program called the BLENREP REMS

They may also choose to enroll in the patient support program called the Together with BLENREP program

Enroll online at [BLENPREMS.com](https://www.blenrep.com) or call 1-855-690-9572

BLENREP is only available through the Risk Evaluation and Management Strategy (REMS) program. Your loved one's healthcare provider will explain what it is, review the Patient Guide, and help them enroll. If your loved one chooses to enroll in Together with BLENREP, they can access additional resources, Nurse Navigators,* and much more. For more information, see pages 12 and 13 of this guide.

Enroll online at [TogetherwithBLENREP.com](https://www.togetherwithblenrep.com)

*The Nurse Navigator does not provide medical advice and is not a part of the doctor's care team. Please ask the doctor any questions that you might have about your loved one's disease and treatment with BLENREP.

2. SCHEDULE EYE APPOINTMENTS

Help your loved one find an eye doctor and schedule eye exams

Your loved one will need to schedule eye exams on a regular basis. It's important to show up for these appointments because delays can push their doses back.

Two eye exams, given at the same eye appointment, are required before treatment starts, before each dose, and as needed for new or worsening symptoms. Your loved one's Nurse Navigator or the oncologist's office can help coordinate.

If needed, you can always use the "Find an Eye Doctor" tool at [BLENREP.com](https://www.blenrep.com)



Slit Lamp Exam



Snellen Eye Chart Exam



These exams do not typically require dilation (widening of the pupil). But the doctor may perform a full examination.

Important Safety Information for BLENREP (cont'd)

Because of the risk of eye problems, BLENREP is available only through a restricted program called the BLENREP Risk Evaluation and Mitigation Strategy (REMS). Your healthcare provider will give you the BLENREP REMS Patient Guide, which describes the risk of eye problems, and will explain the REMS program to you. To receive BLENREP, you must enroll in the REMS program and receive eye exams during treatment. See **"What are the possible side effects of BLENREP?"** for more information about side effects.

3. SCHEDULE ADDITIONAL TESTS


Make sure your loved one is ready for their infusion

Your loved one's doctor may perform additional tests, including checking their blood cell counts, before they start treatment with BLENREP and during treatment as needed.

4. SCHEDULE BLENREP INFUSION

Support your loved one in scheduling the BLENREP infusion (after eye exam results have been shared with the oncologist)

Your loved one's Nurse Navigator or oncologist's office can help schedule the infusion. Keep a calendar of infusion dates and adjust if needed.



Important Safety Information for BLENREP (cont'd)

What are the possible side effects of BLENREP?

BLENREP can cause serious side effects, including:

- See **"Eye Problems"** in **"What is the most important information I should know about BLENREP?"**
- **Decrease in platelets (thrombocytopenia)** is common with BLENREP and can also be severe. Platelets are a type of blood cell that help your blood to clot. Your healthcare provider will check your blood cell counts before you start treatment with BLENREP and during treatment as needed. Tell your healthcare provider if you get any bleeding or bruising during treatment with BLENREP

Please see additional Important Safety Information throughout. Click to see the full [Prescribing Information](#), including [Boxed Warning](#) and [Medication Guide](#) for BLENREP.

BLENREP
belantamab
mafodotin-blmf
for injection 70 mg

Overview of the BLENREP DREAMM-7 study

- The DREAMM-7 study was a global clinical trial. It compared treatment with BLENREP + bortezomib + dexamethasone to a common treatment combination with daratumumab + bortezomib + dexamethasone
- The efficacy of BLENREP was established in a clinical study in adults with multiple myeloma who had received 2 or more prior treatments, including a proteasome inhibitor and an immunomodulatory agent, and experienced disease progression
- Patients were split into 2 groups:
 - 108 people were given **BLENREP + bortezomib + dexamethasone**
 - 109 people were given **daratumumab + bortezomib + dexamethasone**



Helpful terms to know

- **Median:** the middle number in a set of data, where half of the people had longer results than this middle number and half had shorter
- **Progression-free survival:** the length of time people are on treatment before their cancer starts growing and spreading
- **Overall survival:** the length of time people were still alive from when they were randomly assigned to a treatment group
- **Overall response rate:** the percentage of people who respond to a treatment with a partial response or better
- **Remission:** refers to median progression-free survival, the length of time after the start of treatment in which a person is alive and their cancer does not get worse
- **Stringent complete response:** the highest response when cancer cells cannot be found in the bone marrow*
- **Complete response:** when blood and urine tests show no signs of cancer cells*
- **Very good partial response:** when at least 90% of cancer cells have left the blood and urine
- **Partial response:** when cancer cells decrease in the blood and urine by at least 50%



*This does not mean the disease has been cured.

Important Safety Information for BLENREP (cont'd)

The most common side effects of BLENREP when given with bortezomib and dexamethasone include:

- See “Eye problems” in “What is the most important information I should know about BLENREP?”
- upper respiratory tract infection
- liver problems
- diarrhea
- tiredness
- pneumonia
- cataract
- COVID-19

The BLENREP combination delivered proven results



Individual results may vary.

Important Safety Information for BLENREP (cont'd)

The most common severe abnormal blood test results during treatment with BLENREP include decreases in platelets, white blood cells, and hemoglobin, and increases in certain liver enzymes.

Please see additional Important Safety Information throughout. Click to see the full Prescribing Information, including Boxed Warning and Medication Guide for BLENREP.



Eye and vision changes were common with the BLENREP combination and could be severe, but improvement of these changes was possible

89% of those treated with the BLENREP combination had a reduction in vision in at least 1 eye in the BLENREP clinical study, meaning their vision was less sharp, even with corrective lenses

- Changes in vision were measured by the Best-Corrected Visual Acuity (BCVA) test
- Nearly 30% of patients did not experience a change in vision up to 20/50 (see the slight blurring pictured below)



Images are for illustrative purposes only. Vision will differ for each individual. Images created with an Eye Care Professional.

Of people with reduced vision in at least 1 eye, some experienced resolution of their last vision reduction to baseline or better




Important Safety Information for BLENREP (cont'd)

Tell your healthcare provider right away if you get new or worsening unexplained signs or symptoms of lung problems, including shortness of breath, chest pain, or cough. Your healthcare provider may decrease your dose, temporarily stop, or completely stop treatment with BLENREP if you get serious side effects.

BLENREP can affect the cornea—the outermost layer of the eye

Changes occur in the cornea, the clear outer layer at the front of your eye—this is at the surface of the eye, not deep inside.



Supporting your loved one through eye care while on BLENREP

- Help ensure your loved one keeps all eye care appointments because some changes can happen without symptoms and may only be seen on an eye exam
- Ensure your loved one administers their preservative-free artificial tears at least 4 times per day starting with their first infusion and continuing until the end of treatment with BLENREP
 - Your loved one may be eligible for eye drops at no cost through the BLENREP Eye Drop Program. [Click here](#) for more information
- Advise your loved one to avoid wearing contact lenses during treatment unless directed by their eye care specialist
- Make sure your loved one uses caution when driving or operating machinery because BLENREP may cause changes to their vision

Dose changes may help manage eye and vision changes

Based on eye exam results from the eye doctor, your loved one’s oncologist may choose to:



CONTINUE the dose and schedule of BLENREP



REDUCE the dose of BLENREP



PAUSE the dose of BLENREP



STOP BLENREP treatment

Important Safety Information for BLENREP (cont'd)

BLENREP may affect fertility in males and females, which may affect your ability to have children.

Please see additional Important Safety Information throughout. Click to see the full Prescribing Information, including Boxed Warning and Medication Guide for BLENREP.



Getting started with BLENREP

BLENREP infusion



BLENREP will be given at a certified BLENREP REMS outpatient site



BLENREP is given as an intravenous (IV) infusion usually every 3 weeks, in combination with bortezomib and dexamethasone—a healthcare provider will decide how many treatments are needed

If your loved one misses any appointments, call their healthcare provider as soon as possible to reschedule.

BLENREP is a treatment designed to fit into your life



Infusion can be as short as 30 minutes



No hospitalization is needed to start BLENREP



BLENREP may be given by your loved one’s current care team in their community if they participate in REMS



No pre-medications are needed in order to receive BLENREP

Important Safety Information for BLENREP (cont’d)

Talk to your healthcare provider if this is a concern for you.

These are not all of the possible side effects of BLENREP.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

A few helpful reminders for care partners

Review educational materials together.

Read all materials provided by the care team so you know what to expect with treatment.

Attend appointments and stay connected with the care team.

If possible, join your loved one at their appointments, take notes, and ask questions.

Plan eye appointments in advance.

Help your loved one schedule visits and make sure they are using preservative-free artificial tears at least 4 times per day.

Watch for side effects. Monitor for any new or worsening side effects or vision changes, and ensure your loved one reports them to the healthcare team right away.

Remember to care for yourself, too. Be sure to set aside time for rest and self-care whenever possible. There are a few organizations to reach out to for support*:

- [Blood Cancer United](#)
- [Cancer Care](#)
- [Cancer Support Community](#)
- [HealthTree](#)
- [International Myeloma Foundation](#)
- [Multiple Myeloma Research Foundation](#)

*These resources are external to GSK. GSK does not have control over the content or information provided through these resources, and accordingly, does not warrant their accuracy or completeness. These organizations do not endorse any products or services.

“We’re well-informed well before we even go to the doctor.”

–Real care partner like you



Not an actual patient or care partner.

Important Safety Information for BLENREP (cont’d)

Before receiving BLENREP, tell your healthcare provider about all of your medical conditions, including if you:

- have or had vision or eye problems
- have or had bleeding problems
- are pregnant or plan to become pregnant. BLENREP can harm your unborn baby

Please see additional Important Safety Information throughout. Click to see the full [Prescribing Information](#), including [Boxed Warning](#) and [Medication Guide](#) for BLENREP.



The Together with BLENREP program is support made just for your loved one

Here’s what to expect from Together with BLENREP

It’s more than a program—it’s our promise to understand your loved one’s needs, listen to what’s important to them, and create meaningful connections along the way.

Once a doctor has prescribed BLENREP, we will help your loved one to understand the medication coverage process, investigate affordable options, and guide them through moments that matter in their care.



Coverage Support*

Once you help your loved one sign up for Together with BLENREP, their dedicated Nurse Navigator[†] will have their insurance coverage checked and help them understand their benefits. The information provided by Together with BLENREP is not a guarantee of coverage or payment

- Coverage verification for eye exams, infusions, and BLENREP prescriptions



Cost Support[‡]

Along with your loved one’s existing insurance, their dedicated Nurse Navigator can provide information on additional programs that may help pay for BLENREP

- There are options for copay assistance for eye exams and eye drops (subject to eligibility)



Patient Support

Once you help your loved one sign up for Together with BLENREP, a dedicated Nurse Navigator can help them, the oncologist, the Eye Care Professional, and their office staff work together during the treatment journey with BLENREP

*There may be current coverage restrictions or requirements that may apply; be sure to contact your loved one’s plan for more information. The information provided by Together with BLENREP is not a guarantee of coverage.

[†]The Nurse Navigator does not provide medical advice and is not a part of the doctor’s care team. Please ask the doctor any questions that you might have about your loved one’s disease and treatment with BLENREP.

[‡]Together with BLENREP cannot guarantee that your loved one will be eligible for assistance from a foundation, as each foundation has their own rules on eligibility which are subject to change.

Navigating access together: personalized support for coverage, costs, and your loved one’s care journey



You can help your loved one enroll in Together with BLENREP in 2 ways:

Online enrollment

You can start the enrollment process directly online at the **Together with BLENREP website**. You will need to fill out patient information, sign the form digitally, and submit it.

OR

Fax/Paper enrollment

You can **download** and print this form, complete it with the help of an oncologist, and have the office submit it.

To learn more about Together with BLENREP and enroll, [click here](#)

Please see Important Safety Information throughout. Click to see the full [Prescribing Information](#), including [Boxed Warning](#) and [Medication Guide](#) for BLENREP.



Important Safety Information for BLENREP

What is the most important information I should know about BLENREP?

BLENREP can cause serious side effects, including:

- **Eye problems.** Eye problems are common with BLENREP and can also be severe. BLENREP can cause changes to the surface of your eye, which can lead to symptoms of eye problems, including:
 - decreased vision
 - blurred vision
 - dry eyes
 - sensitivity to light
 - feeling like something is in your eyes
 - eye irritation
 - eye pain

Ulcers on the surface of the eye (corneal ulcers), including with infection, may also happen during treatment with BLENREP.

Tell your healthcare provider right away if you notice any new or worsening eye symptoms or vision changes during treatment with BLENREP.

Your healthcare provider will refer you to an eye care specialist (such as an ophthalmologist or optometrist) to check your eyes before you start treatment, before you receive each dose of BLENREP, and as needed for any new or worsening eye problems. **It is important that you:**

- Keep all of your eye care appointments because some changes can happen without symptoms and may only be seen on an eye exam
- Use preservative-free artificial tears at least 4 times per day starting with your first infusion and continuing until the end of treatment with BLENREP
- Avoid wearing contact lenses during treatment with BLENREP unless directed by your eye care specialist
- Use caution when driving or operating machinery because BLENREP may cause changes to your vision

Because of the risk of eye problems, BLENREP is available only through a restricted program called the BLENREP Risk Evaluation and Mitigation Strategy (REMS). Your healthcare provider will give you the BLENREP REMS Patient Guide, which describes the risk of eye problems, and will explain the REMS program to you. To receive BLENREP, you must enroll in the REMS program and receive eye exams during treatment. See **“What are the possible side effects of BLENREP?”** for more information about side effects.

What are the possible side effects of BLENREP?

BLENREP can cause serious side effects, including:

- See **“Eye Problems”** in **“What is the most important information I should know about BLENREP?”**
- **Decrease in platelets (thrombocytopenia)** is common with BLENREP and can also be severe. Platelets are a type of blood cell that help your blood to clot. Your healthcare provider will check your blood cell counts before you start treatment with BLENREP and during treatment as needed. Tell your healthcare provider if you get any bleeding or bruising during treatment with BLENREP

The most common side effects of BLENREP when given with bortezomib and dexamethasone include:

- See **“Eye problems”** in **“What is the most important information I should know about BLENREP?”**
- upper respiratory tract infection
- liver problems
- diarrhea
- tiredness
- pneumonia
- cataract
- COVID-19

The most common severe abnormal blood test results during treatment with BLENREP include decreases in platelets, white blood cells, and hemoglobin, and increases in certain liver enzymes.

Tell your healthcare provider right away if you get new or worsening unexplained signs or symptoms of lung problems, including shortness of breath, chest pain, or cough.

Your healthcare provider may decrease your dose, temporarily stop, or completely stop treatment with BLENREP if you get serious side effects.

BLENREP may affect fertility in males and females, which may affect your ability to have children. Talk to your healthcare provider if this is a concern for you.

These are not all of the possible side effects of BLENREP.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Before receiving BLENREP, tell your healthcare provider about all of your medical conditions, including if you:

- have or had vision or eye problems
- have or had bleeding problems
- are pregnant or plan to become pregnant. BLENREP can harm your unborn baby

Females who are able to become pregnant:

- Your healthcare provider will do a pregnancy test before you start treatment with BLENREP
- Use effective birth control during treatment with BLENREP and for 4 months after the last dose. Talk to your healthcare provider about birth control methods you can use during this time
- Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment with BLENREP

Males with female partners who are able to become pregnant:

- Use effective birth control during treatment with BLENREP and for 6 months after the last dose
- are breastfeeding or plan to breastfeed. It is not known if BLENREP passes into your breast milk. Do not breastfeed during treatment with BLENREP and for 3 months after the last dose

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

You are encouraged to report negative side effects of prescription drugs to the FDA.

Visit www.fda.gov/medwatch or call 1-800-FDA-1088. You may also report negative side effects to GSK at gsk.public.reportum.com or 1-888-825-5249.

Please click to see the full [Prescribing Information](#), including [Boxed Warning](#) and [Medication Guide](#) for BLENREP.



I want to be a

come back kid

CARE PARTNER GUIDE

Not an actual
patient or
care partner.

HELP SUPPORT YOUR LOVED ONE ON THEIR TREATMENT JOURNEY WITH BLENREP

TO LEARN MORE, VISIT [BLENREP.COM](https://blenrep.com)

What is BLENREP?

BLENREP is a prescription medicine used in combination with the medicines bortezomib and dexamethasone to treat adults with multiple myeloma who:

- have received at least 2 prior treatments, including a proteasome inhibitor and an immunomodulatory agent, and
- their cancer has come back or did not respond to prior treatment.

It is not known if BLENREP is safe and effective in children.

Please see additional Important Safety Information throughout. Click to see the full Prescribing Information, including Boxed Warning and Medication Guide for BLENREP.

Important Safety Information

What is the most important information I should know about BLENREP?

BLENREP can cause serious side effects, including:

- **Eye problems.** Eye problems are common with BLENREP and can also be severe. BLENREP can cause changes to the surface of your eye, which can lead to symptoms of eye problems, including:
 - decreased vision
 - blurred vision
 - dry eyes
 - sensitivity to light
 - feeling like something is in your eyes
 - eye irritation
 - eye pain

Trademarks are owned by or licensed to the GSK group of companies.

GSK

©2026 GSK or licensor.
PMUS-BLMBROC260009 August 2026
Produced in USA.

File Name: Care Partner Brochure



BLENREP
belantamab
mafodotin-blmf
for injection 70 mg