



support

made just for you

together

with



BLENREP
belantamab
mafodotin-blmf
for injection 70 mg

Please see Important Safety Information throughout and accompanying [Prescribing Information](#), including Boxed Warning and Medication Guide for BLENREP.

together
with

 **BLENREP**
belantamab
mafodotin-blmf
for injection

It's more than a program—it's **our**
promise to understand **your needs**,
listen to what matters most to you,
and create meaningful **connections**
along the way.

Once your doctor has prescribed
BLENREP, we will help you to
understand the medication coverage
process, investigate **affordable options**,
and **guide you** through moments that
matter in your care.

Please see Important Safety Information throughout and accompanying [Prescribing Information](#),
including Boxed Warning and Medication Guide for BLENREP.

What is BLENREP?

BLENREP is a prescription medicine used in combination with the medicines bortezomib and dexamethasone to treat adults with multiple myeloma who:

- have received at least 2 prior treatments, including a proteasome inhibitor and an immunomodulatory agent, and
- their cancer has come back or did not respond to prior treatment.

It is not known if BLENREP is safe and effective in children.

IMPORTANT SAFETY INFORMATION

What is the most important information I should know about BLENREP?

BLENREP can cause serious side effects, including:

- **Eye problems.** Eye problems are common with BLENREP and can also be severe. BLENREP can cause changes to the surface of your eye, which can lead to symptoms of eye problems, including:
 - o decreased vision
 - o blurred vision
 - o dry eyes
 - o sensitivity to light
 - o feeling like something is in your eyes
 - o eye irritation
 - o eye pain

Ulcers on the surface of the eye (corneal ulcers), including with infection, may also happen during treatment with BLENREP.

Tell your healthcare provider right away if you notice any new or worsening eye symptoms or vision changes during treatment with BLENREP.

Your healthcare provider will refer you to an eye care specialist (such as an ophthalmologist or optometrist) to check your eyes before you start treatment, before you receive each dose of BLENREP, and as needed for any new or worsening eye problems. **It is important that you:**

- o Keep all of your eye care appointments because some changes can happen without symptoms and may only be seen on an eye exam
- o Use preservative-free artificial tears at least 4 times per day starting with your first infusion and continuing until the end of treatment with BLENREP
- o Avoid wearing contact lenses during treatment with BLENREP unless directed by your eye care specialist
- o Use caution when driving or operating machinery because BLENREP may cause changes to your vision

Because of the risk of eye problems, BLENREP is available only through a restricted program called the BLENREP Risk Evaluation and Mitigation Strategy (REMS). Your healthcare provider will give you the BLENREP REMS Patient Guide, which describes the risk of eye problems, and will explain the REMS program to you. To receive BLENREP, you must enroll in the REMS program and receive eye exams during treatment. See **“What are the possible side effects of BLENREP?”** for more information about side effects.

What are the possible side effects of BLENREP?

BLENREP can cause serious side effects, including:

- See **“Eye Problems”** in **“What is the most important information I should know about BLENREP?”**
- **Decrease in platelets (thrombocytopenia)** is common with BLENREP and can also be severe. Platelets are a type of blood cell that help your blood to clot. Your healthcare provider will check your blood cell counts before you start treatment with BLENREP and during treatment as needed. Tell your healthcare provider if you get any bleeding or bruising during treatment with BLENREP

IMPORTANT SAFETY INFORMATION (cont'd)

What are the possible side effects of BLENREP? (cont'd)

The most common side effects of BLENREP when given with bortezomib and dexamethasone include:

- See “**Eye problems**” in “**What is the most important information I should know about BLENREP?**”
- upper respiratory tract infection
- liver problems
- diarrhea
- tiredness
- pneumonia
- cataract
- COVID-19

The most common severe abnormal blood test results during treatment with BLENREP include decreases in platelets, white blood cells, and hemoglobin, and increases in certain liver enzymes.

Tell your healthcare provider right away if you get new or worsening unexplained signs or symptoms of lung problems, including shortness of breath, chest pain, or cough.

Your healthcare provider may decrease your dose, temporarily stop, or completely stop treatment with BLENREP if you get serious side effects.

BLENREP may affect fertility in males and females, which may affect your ability to have children. Talk to your healthcare provider if this is a concern for you. These are not all of the possible side effects of BLENREP.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Before receiving BLENREP, tell your healthcare provider about all of your medical conditions, including if you:

- have or had vision or eye problems
- have or had bleeding problems
- are pregnant or plan to become pregnant. BLENREP can harm your unborn baby

Females who are able to become pregnant:

- o Your healthcare provider will do a pregnancy test before you start treatment with BLENREP
- o Use effective birth control during treatment with BLENREP and for 4 months after the last dose. Talk to your healthcare provider about birth control methods you can use during this time
- o Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment with BLENREP

Males with female partners who are able to become pregnant:

- o Use effective birth control during treatment with BLENREP and for 6 months after the last dose
- are breastfeeding or plan to breastfeed. It is not known if BLENREP passes into your breast milk. Do not breastfeed during treatment with BLENREP and for 3 months after the last dose

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088. You may also report negative side effects to GSK at gsk.public.reportum.com or 1-888-825-5249.

Here's what to expect from Together with BLENREP (belantamab mafodotin-blmf)



Coverage support

We'll check your insurance coverage and help you understand your benefits.*

*The information provided by Together with BLENREP is not a guarantee of coverage or payment.



Cost support

Your dedicated Nurse Navigator can give you information on additional programs that may help you pay for BLENREP.



Patient support

Your Nurse Navigator is at the heart of our program, offering tailored support throughout your treatment journey.

Please see Important Safety Information throughout and accompanying Prescribing Information, including Boxed Warning and Medication Guide for BLENREP.

together
with

BLENREP
belantamab
mafodotin-blmf
for injection

Here's what to expect from your Nurse Navigator[†]



[†]The Nurse Navigator does not provide medical advice and is not a part of your doctor's care team. Please ask your doctor any questions that you might have about your disease and treatment with BLENREP.

[‡]Independent foundations are not affiliated with GSK and are operated independently. Together with BLENREP cannot guarantee that you will be eligible for assistance.

Please see Important Safety Information throughout and accompanying Prescribing Information, including Boxed Warning and Medication Guide for BLENREP.

together
with

BLENREP
belantamab
mafodotin-blmf
for injection

Your dedicated Nurse Navigator can help you with a wealth of topics connected to your treatment journey with BLENREP (belantamab mafodotin-blmf). Not sure about coverage? Don't know where to find financial assistance? Sign up today and get answers from your Nurse Navigator.

Your Nurse Navigator will help provide:



Answers to coverage and benefits questions



Options to help with medication costs



Ongoing support and education about your treatment

Enroll today at
TogetherwithBLENREP.com

Together with BLENREP offers a range of assistance options at no cost to you. First, you and your doctor need to follow these **three easy steps** to complete enrollment.



Step
one

Fill out the form

Select 'ENROLL NOW' to begin your online enrollment.



Step
two

Sign

Once you've completed the form, sign in the appropriate section.



Step
three

Confirm

Once you've completed the form, it will be sent to your doctor.

Questions? Call

1-844-4GSK-ONC (844-447-5662)
Monday–Friday (8:00 AM – 8:00 PM ET)



Eye care support
services:

See the services available to you once you enroll in Together with BLENREP



Help finding an eye care provider in your plan if you don't already have one



Coverage verification for eye exams, infusions, and BLENREP prescriptions



Options for copay assistance for eye exams and eye drops (subject to eligibility)



Eye care appointment reminders

Please see Important Safety Information throughout and accompanying Prescribing Information, including Boxed Warning and Medication Guide for BLENREP.



Connect with us.

1-844-4GSK-ONC (844-447-5662)
Monday–Friday (8:00 AM – 8:00 PM ET)

Visit **TogetherwithGSK.com** for more details
or scan the QR code below to get started.



together
with

 **BLENREP**
belantamab
mafodotin-blmf
for injection

Please see Important Safety Information throughout and accompanying Prescribing Information, including Boxed Warning and Medication Guide for BLENREP.

Trademarks are owned by or licensed to the GSK
group of companies.

©2025 GSK or licensor.

PMUS-BLMCRD250004 October 2025

Produced in USA. 0002-0037-05

GSK