

The BLENREP Eye Drop Program



This brochure gives you key information about the BLENREP Eye Drop Program, which offers eye drops free of charge for patients who are given BLENREP.

Approved Use

BLENREP is a prescription medicine used in combination with the medicines bortezomib and dexamethasone to treat adults with multiple myeloma who:

- have received at least 2 prior treatments, including a proteasome inhibitor and an immunomodulatory agent, and
- their cancer has come back or did not respond to prior treatment

It is not known if BLENREP is safe and effective in children.

Important Safety Information

What is the most important information I should know about BLENREP?

BLENREP can cause serious side effects, including:

- **Eye problems.** Eye problems are common with BLENREP and can also be severe. BLENREP can cause changes to the surface of your eye, which can lead to symptoms of eye problems, including:
 - decreased vision
 - dry eyes
 - blurred vision
 - sensitivity to light
 - feeling like something is in your eyes
 - eye irritation
 - eye pain

Please see additional Important Safety Information throughout. Click to see the full [Prescribing Information](#), including [Boxed Warning and Medication Guide](#) for BLENREP.



BLENREP
belantamab
mafodotin-blmf
for injection 70 mg

Join the BLENREP Eye Drop Program



The Eye Drop Program supports you by sending you eye drops during treatment.

Starting with your first BLENREP infusion, you'll need to use preservative-free, lubricant eye drops at least 4 times a day, or as directed by your doctor. These drops are an important part of your care plan, and the Eye Drop Program is our way of supporting you while on your treatment journey. The drops will be shipped directly to your home for as long as you're on treatment.

Important Safety Information for BLENREP (cont'd)

What is the most important information I should know about BLENREP? (cont'd)

BLENREP can cause serious side effects, including:

Ulcers on the surface of the eye (corneal ulcers), including with infection, may also happen during treatment with BLENREP.

Tell your healthcare provider right away if you notice any new or worsening eye symptoms or vision changes during treatment with BLENREP.

Your healthcare provider will refer you to an eye care specialist (such as an ophthalmologist or optometrist) to check your eyes before you start treatment, before you receive each dose of BLENREP, and as needed for any new or worsening eye problems. **It is important that you:**

- Keep all of your eye care appointments because some changes can happen without symptoms and may only be seen on an eye exam
- Use preservative-free artificial tears at least 4 times per day starting with your first infusion and continuing until the end of treatment with BLENREP
- Avoid wearing contact lenses during treatment with BLENREP unless directed by your eye care specialist
- Use caution when driving or operating machinery because BLENREP may cause changes to your vision

Frequently asked questions

Eligibility

Q. How do I know if I'm eligible for the Eye Drop Program?

- When your doctor prescribes BLENREP, you will need to enroll in the BLENREP Risk Evaluation and Mitigation Strategy (REMS) program
- You must be a resident of the US (including the District of Columbia, Puerto Rico, and the US Virgin Islands)
- Your doctor prescribing BLENREP must also be a resident of the US
- You can enroll in the eye drop program by checking the box on the Together with BLENREP enrollment form if you want eye drops. Enrollment in Together with BLENREP is not required in order to get eye drops

Enrollment

Q. How do I enroll in the program?

To enroll, you and your doctor should complete applicable sections of the Together with BLENREP enrollment form. Be sure to mark the box next to the Eye Drop Program when you complete the form available on the Together with BLENREP website.


Q. After I enroll, how will I receive my eye drops?

After you enroll, the Eye Drop Program will reach out to you. The Eye Drop Program will confirm where to ship the eye drops so they go directly to your home.

Important Safety Information for BLENREP (cont'd)

Because of the risk of eye problems, BLENREP is available only through a restricted program called the BLENREP Risk Evaluation and Mitigation Strategy (REMS). Your healthcare provider will give you the BLENREP REMS Patient Guide, which describes the risk of eye problems, and will explain the REMS program to you. To receive BLENREP, you must enroll in the REMS program and receive eye exams during treatment. See **“What are the possible side effects of BLENREP?”** for more information about side effects.

Please see additional Important Safety Information throughout. Click to see the full [Prescribing Information](#), including [Boxed Warning](#) and [Medication Guide](#) for BLENREP.



Frequently asked questions (cont'd)



About the eye drops

Q. Can I choose my own brand of eye drops?

You can use other eye drops if you wish; however, they won't be part of the no-cost program. If you do purchase your own eye drops, it's important that you choose drops that are preservative free. You should follow the advice of your eye care professional. BLENREP does not endorse any specific brand.

Q. Who can answer questions about the Eye Drop Program or the preservative-free eye drops?

You can reach out to the Eye Drop Program for any questions. They are there to help. Call them at 1-844-4GSK-ONC (1-844-447-5662), Monday-Friday, 8:00 AM-8:00 PM ET.

Q. How long will my eye drops last?

After we confirm your BLENREP prescription and your enrollment in REMS, you'll receive an initial shipment with a 60-vial supply of eye drops. Eye drops will be shipped to your address.

After your continued use of BLENREP and your enrollment in REMS is confirmed, you'll receive an additional 60-vial supply of eye drops every 50 days for the next 2-month period. This will continue until you stop treatment or opt out of the program. The Eye Drop Program will check on day 47 to confirm you're still on treatment.

Important Safety Information for BLENREP (cont'd)

What are the possible side effects of BLENREP?

BLENREP can cause serious side effects, including:

- See "Eye Problems" in "What is the most important information I should know about BLENREP?"
- **Decrease in platelets (thrombocytopenia)** is common with BLENREP and can also be severe. Platelets are a type of blood cell that help your blood to clot. Your healthcare provider will check your blood cell counts before you start treatment with BLENREP and during treatment as needed. Tell your healthcare provider if you get any bleeding or bruising during treatment with BLENREP

Frequently asked questions (cont'd)

About the eye drops (cont'd)

Q. How should I store my eye drops?

The eye drops should be stored at room temperature.

Q. Can I use the eye drops with my contact lenses?

No, contact lenses for correcting your vision should not be worn while taking BLENREP, unless directed by your doctor.

Cost/insurance

Q. Is there any cost associated with the program?

No, there is no cost associated with this program for eligible patients.

Q. Do I need to provide insurance information to take part in the Eye Drop Program?

No, you don't need to provide any insurance information if you only want to take part in the Eye Drop Program. On the Together with BLENREP enrollment form, all you need to do is check the box on the enrollment form that you want to enroll along with providing a signature from you and your doctor.

Important Safety Information for BLENREP (cont'd)


The most common side effects of BLENREP when given with bortezomib and dexamethasone include:

- See "Eye problems" in "What is the most important information I should know about BLENREP?"
- diarrhea
- upper respiratory tract infection
- liver problems
- tiredness
- pneumonia
- cataract
- COVID-19

The most common severe abnormal blood test results during treatment with BLENREP include decreases in platelets, white blood cells, and hemoglobin, and increases in certain liver enzymes.

Tell your healthcare provider right away if you get new or worsening unexplained signs or symptoms of lung problems, including shortness of breath, chest pain, or cough.

Please see additional Important Safety Information throughout. Click to see the full [Prescribing Information](#), including [Boxed Warning](#) and [Medication Guide](#) for BLENREP.

 **BLENREP**
belantamab
mafodotin-blmf
for injection 70 mg

Important Safety Information for BLENREP

What is the most important information I should know about BLENREP?

BLENREP can cause serious side effects, including:

- **Eye problems.** Eye problems are common with BLENREP and can also be severe. BLENREP can cause changes to the surface of your eye, which can lead to symptoms of eye problems, including:
 - decreased vision
 - blurred vision
 - dry eyes
 - sensitivity to light
 - feeling like something is in your eyes
 - eye irritation
 - eye pain

Ulcers on the surface of the eye (corneal ulcers), including with infection, may also happen during treatment with BLENREP.

Tell your healthcare provider right away if you notice any new or worsening eye symptoms or vision changes during treatment with BLENREP.

Your healthcare provider will refer you to an eye care specialist (such as an ophthalmologist or optometrist) to check your eyes before you start treatment, before you receive each dose of BLENREP, and as needed for any new or worsening eye problems. **It is important that you:**

- Keep all of your eye care appointments because some changes can happen without symptoms and may only be seen on an eye exam
- Use preservative-free artificial tears at least 4 times per day starting with your first infusion and continuing until the end of treatment with BLENREP
- Avoid wearing contact lenses during treatment with BLENREP unless directed by your eye care specialist
- Use caution when driving or operating machinery because BLENREP may cause changes to your vision

Because of the risk of eye problems, BLENREP is available only through a restricted program called the BLENREP Risk Evaluation and Mitigation Strategy (REMS). Your healthcare provider will give you the BLENREP REMS Patient Guide, which describes the risk of eye problems, and will explain the REMS program to you. To receive BLENREP, you must enroll in the REMS program and receive eye exams during treatment. See **“What are the possible side effects of BLENREP?”** for more information about side effects.

What are the possible side effects of BLENREP?

BLENREP can cause serious side effects, including:

- See **“Eye Problems”** in **“What is the most important information I should know about BLENREP?”**
- **Decrease in platelets (thrombocytopenia)** is common with BLENREP and can also be severe. Platelets are a type of blood cell that help your blood to clot. Your healthcare provider will check your blood cell counts before you start treatment with BLENREP and during treatment as needed. Tell your healthcare provider if you get any bleeding or bruising during treatment with BLENREP

The most common side effects of BLENREP when given with bortezomib and dexamethasone include:

- See **“Eye problems”** in **“What is the most important information I should know about BLENREP?”**
- upper respiratory tract infection
- liver problems
- diarrhea
- tiredness
- pneumonia
- cataract
- COVID-19

The most common severe abnormal blood test results during treatment with BLENREP include decreases in platelets, white blood cells, and hemoglobin, and increases in certain liver enzymes.

Tell your healthcare provider right away if you get new or worsening unexplained signs or symptoms of lung problems, including shortness of breath, chest pain, or cough.

Your healthcare provider may decrease your dose, temporarily stop, or completely stop treatment with BLENREP if you get serious side effects.

BLENREP may affect fertility in males and females, which may affect your ability to have children. Talk to your healthcare provider if this is a concern for you.

These are not all of the possible side effects of BLENREP.

Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.

Before receiving BLENREP, tell your healthcare provider about all of your medical conditions, including if you:

- have or had vision or eye problems
- have or had bleeding problems
- are pregnant or plan to become pregnant. BLENREP can harm your unborn baby

Females who are able to become pregnant:

- Your healthcare provider will do a pregnancy test before you start treatment with BLENREP
- Use effective birth control during treatment with BLENREP and for 4 months after the last dose. Talk to your healthcare provider about birth control methods you can use during this time
- Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment with BLENREP

Males with female partners who are able to become pregnant:

- Use effective birth control during treatment with BLENREP and for 6 months after the last dose
- are breastfeeding or plan to breastfeed. It is not known if BLENREP passes into your breast milk. Do not breastfeed during treatment with BLENREP and for 3 months after the last dose

Tell your healthcare provider about all the medicines you take, including prescription and over-the-counter medicines, vitamins, and herbal supplements.

You are encouraged to report negative side effects of prescription drugs to the FDA. Visit www.fda.gov/medwatch or call 1-800-FDA-1088. You may also report negative side effects to GSK at gsk.public.reportum.com or 1-888-825-5249.



Please click to see the full [Prescribing Information](#), including [Boxed Warning](#) and [Medication Guide](#) for BLENREP.

GSK supports you on your treatment journey with eye drops delivered to your home



Click the link below to enroll in the **BLENREP Eye Drop Program** and begin receiving preservative-free, lubricant eye drops.



We are here to help!

You can reach the Eye Drop Program at **1-844-4GSK-ONC** (1-844-447-5662)

Monday–Friday 8:00 AM–8:00 PM ET
Or visit togetherwithBLENREP.com

Please see Important Safety Information throughout. Click to see the full Prescribing Information, including Boxed Warning and Medication Guide for BLENREP.

Trademarks are owned by or licensed to the GSK group of companies.



©2025 GSK or licensor.
PMUS-BLMLBND250042 December 2025
Produced in USA.

File Name: Eye Drop Program



BLENREP
belantamab
mafodotin-blmf
for injection 70 mg