

# Partnering with your care team



**This guide will introduce you to the range of individuals who may be part of your care team. This support network includes nurses, oncologists, ophthalmologists, and optometrists who will work with you throughout your treatment journey.**

## Approved Use

BLENREP is a prescription medicine used in combination with the medicines bortezomib and dexamethasone to treat adults with multiple myeloma who:

- have received at least 2 prior treatments, including a proteasome inhibitor and an immunomodulatory agent, and
- their cancer has come back or did not respond to prior treatment

It is not known if BLENREP is safe and effective in children.

## Important Safety Information

**What is the most important information I should know about BLENREP?**

**BLENREP can cause serious side effects, including:**

- **Eye problems.** Eye problems are common with BLENREP and can also be severe. BLENREP can cause changes to the surface of your eye, which can lead to symptoms of eye problems, including:
  - decreased vision
  - dry eyes
  - feeling like something is in your eyes
  - eye irritation
  - blurred vision
  - sensitivity to light
  - eye pain

Please see additional Important Safety Information throughout. Click to see the full [Prescribing Information, including Boxed Warning and Medication Guide for BLENREP.](#)



**BLENREP**  
belantamab  
mafodotin-blmf  
for injection 70 mg

## Meet your care team

While your oncologist is the main person who oversees your care, there is a whole team of specialists working with you and your doctor to help you through your treatment journey.



### The following people may become part of your care team throughout your BLENREP journey



#### Oncology Nurses

specialize in cancer care. Your oncology nurse can provide multiple myeloma education and coordinate treatment plans with oncologists. They can also talk to you about side effects and what to expect when on treatment.



#### Ophthalmologists and Optometrists

perform eye exams before each dose of BLENREP. Additional eye exams will also be needed for new or worsening symptoms or as your doctor recommends. You can visit either of these eye care professionals for the eye exams you'll need.

### The following people may become part of your care team throughout your BLENREP journey (cont'd)



#### Hematologists and Hematology Nurses

specialize in blood cancers like multiple myeloma. They work closely with your oncologist to help manage your treatment and monitor lab results.



#### Infusion Nurses

administer intravenous (IV) treatments like chemotherapy and immunotherapy. They will be the ones administering your treatment with BLENREP. Your infusion nurse will monitor for side effects and ensure safety during infusions.

### Important Safety Information for BLENREP (cont'd)

Ulcers on the surface of the eye (corneal ulcers), including with infection, may also happen during treatment with BLENREP.

**Tell your healthcare provider right away** if you notice any new or worsening eye symptoms or vision changes during treatment with BLENREP.

Your healthcare provider will refer you to an eye care specialist (such as an ophthalmologist or optometrist) to check your eyes before you start treatment, before you receive each dose of BLENREP, and as needed for any new or worsening eye problems. **It is important that you:**

- Keep all of your eye care appointments because some changes can happen without symptoms and may only be seen on an eye exam
- Use preservative-free artificial tears at least 4 times per day starting with your first infusion and continuing until the end of treatment with BLENREP
- Avoid wearing contact lenses during treatment with BLENREP unless directed by your eye care specialist
- Use caution when driving or operating machinery because BLENREP may cause changes to your vision

### Important Safety Information for BLENREP (cont'd)

**Because of the risk of eye problems, BLENREP is available only through a restricted program called the BLENREP Risk Evaluation and Mitigation Strategy (REMS).**

Your healthcare provider will give you the BLENREP REMS Patient Guide, which describes the risk of eye problems, and will explain the REMS program to you. To receive BLENREP, you must enroll in the REMS program and receive eye exams during treatment. See **“What are the possible side effects of BLENREP?”** for more information about side effects.

**Please see additional Important Safety Information throughout. Click to see the full Prescribing Information, including Boxed Warning and Medication Guide for BLENREP.**

 **BLENREP**  
belantamab  
mafodotin-blmf  
for injection 70 mg

# Navigating care team conversations

Having constructive conversations with your care team can help them know what is on your mind and how best to respond to your needs. See below for tips and possible topics to discuss.



## Tips for having constructive conversations with your care team



### Provide the full picture

- It is important to be open with your care team so they fully understand your experiences. This helps them to better manage your treatment. If you experience any side effects, be sure to share them with your care team



### Keep track of how you feel between visits

- It can be hard to remember what happened in previous weeks. Take a few minutes to write down how you feel each day throughout treatment and bring it with you to appointments



### Bring someone for support

- Consider bringing a friend or loved one to your appointment for support and to help advocate on your behalf, if needed

## Tips for having constructive conversations with your care team (cont'd)



### Share what is important to you

- Talk with your care team to help align your care plan with your goals and anything else that matters to you. You can update your goals as they change and share them with your doctor and nurse
- You can also make a list of questions to ask your care team before your next appointment



### Advocate for yourself to receive the information you need

- When you are with your care team, speak up if you don't understand something to get the information you need. Your care team may not know what you are feeling until you tell them
- In addition, ask your care team about enrolling in Together with BLENREP—a support program offered by the makers of BLENREP for people taking BLENREP. You can see the back cover for additional details

Keep in mind that this list may not cover everyone you'll see in your BLENREP treatment journey.



## Important Safety Information for BLENREP (cont'd)

### What are the possible side effects of BLENREP?

#### BLENREP can cause serious side effects, including:

- See "Eye Problems" in "What is the most important information I should know about BLENREP?"
- **Decrease in platelets (thrombocytopenia)** is common with BLENREP and can also be severe. Platelets are a type of blood cell that help your blood to clot. Your healthcare provider will check your blood cell counts before you start treatment with BLENREP and during treatment as needed. Tell your healthcare provider if you get any bleeding or bruising during treatment with BLENREP

## Important Safety Information for BLENREP (cont'd)

### The most common side effects of BLENREP when given with bortezomib and dexamethasone include:

- See "Eye problems" in "What is the most important information I should know about BLENREP?"
- upper respiratory tract infection
- liver problems
- diarrhea
- tiredness
- pneumonia
- cataract
- COVID-19

Please see additional Important Safety Information throughout. Click to see the full [Prescribing Information](#), including [Boxed Warning](#) and [Medication Guide](#) for BLENREP.

 **BLENREP**  
belantamab  
mafodotin-blmf  
for injection 70 mg

## Important Safety Information for BLENREP

### What is the most important information I should know about BLENREP?

#### BLENREP can cause serious side effects, including:

- **Eye problems.** Eye problems are common with BLENREP and can also be severe. BLENREP can cause changes to the surface of your eye, which can lead to symptoms of eye problems, including:
  - decreased vision
  - blurred vision
  - dry eyes
  - sensitivity to light
  - feeling like something is in your eyes
  - eye irritation
  - eye pain

Ulcers on the surface of the eye (corneal ulcers), including with infection, may also happen during treatment with BLENREP.

**Tell your healthcare provider right away** if you notice any new or worsening eye symptoms or vision changes during treatment with BLENREP.

Your healthcare provider will refer you to an eye care specialist (such as an ophthalmologist or optometrist) to check your eyes before you start treatment, before you receive each dose of BLENREP, and as needed for any new or worsening eye problems. **It is important that you:**

- Keep all of your eye care appointments because some changes can happen without symptoms and may only be seen on an eye exam
- Use preservative-free artificial tears at least 4 times per day starting with your first infusion and continuing until the end of treatment with BLENREP
- Avoid wearing contact lenses during treatment with BLENREP unless directed by your eye care specialist
- Use caution when driving or operating machinery because BLENREP may cause changes to your vision

**Because of the risk of eye problems, BLENREP is available only through a restricted program called the BLENREP Risk Evaluation and Mitigation Strategy (REMS).** Your healthcare provider will give you the BLENREP REMS Patient Guide, which describes the risk of eye problems, and will explain the REMS program to you. To receive BLENREP, you must enroll in the REMS program and receive eye exams during treatment. See **“What are the possible side effects of BLENREP?”** for more information about side effects.

### What are the possible side effects of BLENREP?

#### BLENREP can cause serious side effects, including:

- See **“Eye Problems”** in **“What is the most important information I should know about BLENREP?”**
- **Decrease in platelets (thrombocytopenia)** is common with BLENREP and can also be severe. Platelets are a type of blood cell that help your blood to clot. Your healthcare provider will check your blood cell counts before you start treatment with BLENREP and during treatment as needed. Tell your healthcare provider if you get any bleeding or bruising during treatment with BLENREP

**The most common side effects of BLENREP when given with bortezomib and dexamethasone include:**

- See **“Eye problems”** in **“What is the most important information I should know about BLENREP?”**
- upper respiratory tract infection
- liver problems
- diarrhea
- tiredness
- pneumonia
- cataract
- COVID-19

**The most common severe abnormal blood test results during treatment with BLENREP include** decreases in platelets, white blood cells, and hemoglobin, and increases in certain liver enzymes.

Tell your healthcare provider right away if you get new or worsening unexplained signs or symptoms of lung problems, including shortness of breath, chest pain, or cough.

Your healthcare provider may decrease your dose, temporarily stop, or completely stop treatment with BLENREP if you get serious side effects.

BLENREP may affect fertility in males and females, which may affect your ability to have children. Talk to your healthcare provider if this is a concern for you.

These are not all of the possible side effects of BLENREP.

**Call your doctor for medical advice about side effects. You may report side effects to FDA at 1-800-FDA-1088.**

**Before receiving BLENREP, tell your healthcare provider about all of your medical conditions, including if you:**

- have or had vision or eye problems
- have or had bleeding problems
- are pregnant or plan to become pregnant. BLENREP can harm your unborn baby
  - Females who are able to become pregnant:**
    - Your healthcare provider will do a pregnancy test before you start treatment with BLENREP
    - Use effective birth control during treatment with BLENREP and for 4 months after the last dose. Talk to your healthcare provider about birth control methods you can use during this time
    - Tell your healthcare provider if you become pregnant or think you may be pregnant during treatment with BLENREP
  - Males with female partners who are able to become pregnant:**
    - Use effective birth control during treatment with BLENREP and for 6 months after the last dose
- are breastfeeding or plan to breastfeed. It is not known if BLENREP passes into your breast milk. Do not breastfeed during treatment with BLENREP and for 3 months after the last dose

**Tell your healthcare provider about all the medicines you take,** including prescription and over-the-counter medicines, vitamins, and herbal supplements.

**You are encouraged to report negative side effects of prescription drugs to the FDA. Visit [www.fda.gov/medwatch](http://www.fda.gov/medwatch) or call 1-800-FDA-1088. You may also report negative side effects to GSK at [gsk.public.reportum.com](http://gsk.public.reportum.com) or 1-888-825-5249.**



**Please click to see the full [Prescribing Information](#), including [Boxed Warning](#) and [Medication Guide](#) for BLENREP.**

**If you haven't enrolled in Together with BLENREP, sign up today!**



**Click the link below to enroll and access additional resources, Nurse Navigators,\* and much more.**



**We are here to help!**

You can reach your Nurse Navigator at **1-844-4GSK-ONC** (1-844-447-5662)

Monday–Friday 8:00 AM–8:00 PM ET

Or visit [togetherwithBLENREP.com](https://togetherwithBLENREP.com)

\*The Nurse Navigator does not provide medical advice and is not a part of your doctor's care team. Please ask your doctor any questions that you might have about your disease and treatment with BLENREP.

**Please see Important Safety Information throughout. Click to see the full Prescribing Information, including Boxed Warning and Medication Guide for BLENREP.**

Trademarks are owned by or licensed to the GSK group of companies.



©2025 GSK or licensor.  
PMUS-BLMLBND250040 December 2025  
Produced in USA.

File Name: Partnering With Your Care Team



**BLENREP**  
belantamab  
mafodotin-blmf  
for injection 70 mg